



PIG STARTER

To Increase the Digestibility of Phytin-Bound Phosphorus in Complete Swine Starter Diets

GUARANTEED ANALYSIS:

Crude Protein, min.....	21.0%
Crude Fat, min.....	7.0%
Crude Fiber, max.....	3.5%
Calcium, min.....	0.6%
Calcium, max.....	1.1%
Phosphorus, min.....	0.7%
Salt, min.....	0.3%
Salt, max.....	0.8%
Lysine, min.....	1.5%
Selenium, min.....	0.3 ppm
Zinc, min.....	3,000 ppm
Phytase, min*.....	907 FTU/lb.

*One unit of phytase activity (FTU) is defined as the quantity of enzyme which liberates 1 micromol of inorganic phosphorus per minute. Environmental factors such as feed pH, moisture and processing methods such as pelleting and extrusion may affect enzyme activity.

FEEDING INSTRUCTIONS:

Feed as a sole ration to starting pigs from 26 to 40 days of age at the rate of 0.5 pound per 10 pounds of body weight. This will provide 10 milligrams Chlortetracycline per pound of body weight per day. A source of water is also necessary.

WARNING: Withdraw 2 days before slaughter.

INGREDIENTS:

Please refer to the product tag/label for the ingredient list.

LIMITATIONS FOR USE: Feed approximately 400 g/ton of Chlortetracycline, varying with bodyweight and feed consumption to provide 10 mg/lb. of bodyweight per day. Feed continuously as sole ration for up to 14 days.

Hubbard does not use "Restricted-use Proteins" in their products and is in compliance with FDA and state requirements regarding the use, handling and storage of "Restricted-use Protein" products.

PROFILE:

Pig Starter is the most highly used and talked-about starter on the market. Whether you are getting your pigs ready for sale or just want them to bloom and grow, Pig Starter works. Designed for pigs 21 days of age and older.

THIS IS A PRODUCT INFORMATION SHEET ONLY IN NO WAY SHOULD THIS BE UTILIZED AS A LABEL

Options in a 50-lb bag #50MM or bulk invoice